

## PLAYING WITH EIGENVALUES-PUTNAM SEMINAR 2016

1. Determine the number of possible values for the determinant of  $A$ , given that  $A$  is a  $n \times n$  matrix with real entries such that  $A^3 - A^2 - 3A + 2I = 0$ , where  $I$  is the identity and  $0$  is the all-zero matrix.
2. Consider matrices  $A_1, A_2, \dots, A_m \in M_n(\mathbb{R})$  not all nilpotent. Prove that there is an integer number  $k > 0$  such that  $A_1^k + A_2^k + \dots + A_m^k \neq O_n$ .
3. Let  $A, B \in M_2(\mathbb{C})$  such that :  $A^2 + B^2 = 2AB$  .
  - a) Prove that :  $AB = BA$  .
  - b) Prove that :  $\text{tr}(A) = \text{tr}(B)$  .
4. Let  $A, B$  be matrices of dimension  $2010 \times 2010$  which commute and have real entries, such that  $A^{2010} = B^{2010} = I$ , where  $I$  is the identity matrix. Prove that if  $\text{tr}(AB) = 2010$ , then  $\text{tr}(A) = \text{tr}(B)$ .
5. Let  $A$  and  $B$  be real symmetric matrixes with all eigenvalues strictly greater than 1. Let  $\lambda$  be a real eigenvalue of matrix  $AB$ . Prove that  $|\lambda| > 1$ .
6. Does there exist a real  $3 \times 3$  matrix  $A$  such that  $\text{tr}(A) = 0$  and  $A^2 + A^t = I$ ? ( $\text{tr}(A)$  denotes the trace of  $A$ ,  $A^t$  the transpose of  $A$ , and  $I$  is the identity matrix.)
7. let  $A, B \in M_n(\mathbb{C})$ . If  $A(AB - BA) = (AB - BA)A$  prove that  $AB - BA$  is nilpotent.
8. Let  $A$  be a real symmetric matrix and  $B \in M_n(\mathbb{C})$  such that  $AB + BA = 0$  . Prove :  $AB = 0$
9. Let  $A \in \text{Sl}_3(\mathbb{Z})$  of finite order. Find all possible values of  $\text{tr}(A)$ .
10. Let  $A$  and  $B$  be two complex matrices. Prove that the following conditions are equivalent:
  - a) For any  $M \in M_n(\mathbb{C})$  the characteristic polynomials of  $AM$  and  $AM + B$  are the same
  - b)  $B$  is nilpotent and  $BA = O_n$ .
11.  $A, B \in M_2(\mathbb{C})$  with  $\det(A) = 1, |\text{tr}(A)| \neq 2, \det(B) = 1, -\text{tr}(B) \neq 2$  and suppose also  $A, B$  do not have common eigenvectors. Given that there exist  $(n_1, \dots, n_k, m_1, \dots, m_k) \in \mathbb{Z}$  such that  $A^{n_1} B^{m_1} \dots A^{n_k} B^{m_k} = I_2$  prove that  $A^{-n_1} B^{-m_1} \dots A^{-n_k} B^{-m_k} = I_2$
12. Let  $A, B \in M_2(\mathbb{C})$  with  $\exp(A) = \exp(B)$ . Suppose for any eigenvalue  $a$  of  $A$  and  $b$  of  $B$ ,  $a - b \notin 2\pi i\mathbb{Z}$ . Then  $A = B$ .
13. Let  $A, B$  be square matrices a) of size  $2016 \times 2016$ ; b) of size  $2017 \times 2017$ . Do there necessarily exist real numbers  $a, b$  such that  $a^2 + b^2 \neq 0$  and the matrix  $aA + bB$  is singular?
14. Let a square matrix  $P$  be neither zero nor unit and such that  $P^2 = P$ . Does there always exist such a matrix  $Q$  that  $Q^2 = Q, PQ = QPQ$  but  $QP \neq PQ$ ?